

N-alkylidene piperidine compounds and their enantiomers, process for their preparation and pharmaceutical compositions containing them.

Publication number: EP0515240

Publication date: 1992-11-25

Inventor: EMONDS-ALT XAVIER (FR); MARTINEZ SERGE (FR); PROIETTO VINCENZO (FR); VAN BROECK DIDIER (FR)

Applicant: SANOFI ELF (FR)

Classification:















- international: **A61K31/44; A61K31/4427; A61K31/4433; A61K31/445; A61K31/4465; A61K31/4468; A61K31/454; A61K31/4545; A61P43/00; C07D211/46; C07D211/54; C07D211/58; C07D401/06; C07D401/12; C07D403/12; C07D409/12; A61K31/44; A61K31/4427; A61K31/445; A61K31/4465; A61K31/4468; A61K31/4523; A61P43/00; C07D211/00; C07D401/00; C07D403/00; C07D409/00; (IPC1-7): A61K31/445; C07D211/46; C07D211/54; C07D211/58; C07D401/12; C07D401/14; C07D409/12**

- European: C07D211/46; C07D211/54; C07D211/58; C07D401/12

Application number: EP19920401237 19920430





Priority number(s): FR19910005486 19910503

Also published as:

 US5411971 (A1)
 JP5140103 (A)
 IE921365 (A1)
 FR2676054 (A1)
 FI921950 (A)
 BR9201655 (A)
 EP0515240 (B1)
 RU2089547 (C1)
 NO178572B (B)
 HU213915 (B)
 FI103041B (B)
 CZ282919 (B6)
 CA2067924 (C)
 AU657321B (B2)

less <<

Cited documents:

 DE2719211
 DE2840302
 DE2610228
 EP0428434

Report a data error here

Abstract of EP0515240

The invention relates to compounds of formula: in which: - m is equal to 2 or 3; - Ar represents a phenyl which is unsubstituted or substituted one or more times by a halogen atom, preferably a chlorine or fluorine atom, by a C1-C3 alkyl, by a trifluoromethyl, by an alkoxy in which the alkyl is C1-C3, by a hydroxyl or by a methylenedioxy; a thienyl, pyridyl or imidazolyl group which is unsubstituted or substituted by a C1-C3 alkyl; - Ar' represents a phenyl group which is unsubstituted or mono- or disubstituted by a halogen atom, preferably a chlorine or fluorine atom, by a C1-C3 alkyl, by a trifluoromethyl, by an alkoxy in which the alkyl is C1-C3, by hydroxyl or by a methylenedioxy; a thienyl group; an imidazolyl group or a benzothienyl group which is unsubstituted or substituted by a halogen, preferably by a chlorine or fluorine atom; a naphthyl group which is unsubstituted or substituted by a halogen, preferably by a fluorine atom; a biphenyl group; an indolyl which is unsubstituted or substituted on the nitrogen by a benzyl group; - X represents an oxygen atom, a sulphur atom, a sulphone or a sulfoxide, an -NH- group, a group or a group in which Alk is a C1-C3 alkyl group or a group in which Alk represents a C1-C3 alkylene and X1 and X2 independently represent hydrogen, a C1-C3 alkyl or constitute, with the nitrogen atom to which they are bonded, a heterocycle chosen from pyrrolidine, piperidine or morpholine; - Q represents hydrogen, a C1-C4 alkyl group or an aminoalkyl group of formula -(CH2)q-Am', where q is 2 or 3 and Am' is a piperidino, 4-benzylpiperidino or dialkylamino group, it being possible for each alkyl to contain from 1 to 4 carbon atoms; - R represents hydrogen, a methyl group or a group (CH2)n-L, where n is an integer from 2 to 6 and L is hydrogen or an amino group; - T represents a group chosen from: W being an oxygen or sulphur atom, and - Z represents either M or OM when T represents: the group or M when T represents the group M represents hydrogen or a straight or branched C1-C6 alkyl; an alpha-hydroxybenzyl, an alpha-alkylbenzyl or a phenylalkyl in which the alkyl group comprises 1 to 3 carbon atoms, unsubstituted or mono- or polysubstituted on the aromatic ring by a halogen, a hydroxyl, an alkoxy containing 1 to 4 carbon atoms, or an alkyl containing 1 to 4 carbon atoms; a pyridylalkyl in which the alkyl group comprises 1 to 3 carbon atoms; a naphthylalkyl in which the alkyl

group comprises 1 to 3 carbon atoms; a pyridylthioalkyl in which the alkyl group comprises 1 to 3 carbon atoms; a styryl; a 1-methylimidazol-2-ylthioalkyl in which the alkyl group comprises 1 to 3 carbon atoms; a 1-oxo-3-phenylindan-2-yl; an unsubstituted, mono- or polysubstituted aromatic or heteroaromatic group; or to one of their salts with inorganic or organic acids or to one of their quaternary ammonium salts formed with the nitrogen of piperidine. Application: Antagonists of receptors of neurokinins (neurokinins).

.....
Data supplied from the **esp@cenet** database - Worldwide